

Graphene-based fine tuning of Fano resonance transmission of quasi-bound states in the continuum: supplement

MYUNGHWAN KIM,^{1,*}  CHUL-SIK KEE,^{2,3}  AND SOEUN KIM^{1,4}

¹*Division of Applied Photonics System Research, Advanced Photonics Research Institute, Gwangju Institute of Science and Technology, Gwangju 61005, Republic of Korea*

²*Division of Frontier Photonics Research, Advanced Photonics Research Institute, Gwangju Institute of Science and Technology, Gwangju 61005, Republic of Korea*

³*cskee@gist.ac.kr*

⁴*sekim@gist.ac.kr*

**kimmh1028@gist.ac.kr*

This supplement published with Optica Publishing Group on 5 August 2022 by The Authors under the terms of the [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/) in the format provided by the authors and unedited. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI.

Supplement DOI: <https://doi.org/10.6084/m9.figshare.20390646>

Parent Article DOI: <https://doi.org/10.1364/OE.468890>

**Graphene-based fine tuning of Fano
resonance transmission of quasi-bound states
in the continuum
: supplemental document**




Fig. S1. Quality factor (Q-factor) as a function of Fermi level.